Art. IX. — CONTRIBUTIONS TO ENCEPHALIC ANATOMY.

By E. C. Spitzka.

PART VIII.

The Brain of the Iguana.

HAVING, through a piece of good fortune, come into the possession of a living iguana, and thence obtained the brain and cord in a perfectly fresh condition, I was enabled to make a study for the first time of the remarkable brain of this saurian.

As regards the exterior of the encephalon, it presents nothing very different from that of any other higher reptile. On a lateral view, however, it exhibits a much acuter basilar incurvation, approximating to the bird's brain in this respect. As in birds, also, the optic nerves leave the skull directly on emerging from the chiasm. It is remarked also that the optic lobes are far larger than in any reptile or bird thus far examined by anatomists, in fact, excluding the case of the finny tribes, it may be said that the iguana possesses the largest optic lobes in the animal kingdom. They are as massive in their grey and white tissues, and nearly as voluminous as the cerebral hemispheres.

The olfactory lobes and bulbs offer nothing special for consideration.

On a transverse section through the cerebral hemispheres, I am able to identify the component parts of the cornu ammonis of the mammalia. It appears that the medial thin wall of the cerebral vesicle corresponds, with its layer of closely packed pyramidal nerve cells, to the *stratum corporum nervorum arctorum* of Kuppfer, and it is indeed separated from the cortical layer of the convexity, which I believe to correspond, as far as the thin part extends, to the *sigma* of the cornu. At the nuclei of the third and fourth pairs constitute a common cell mass, unlike the relation in the mammalia, and that the third and fourth pairs arise almost in the same plane, the third from the ventral, the fourth from the dorsal extensions of the common nucleus.

I would call attention to the fact that the average dimensions of the cell nuclei of the auditory nerve nucleus equal those of the motor nuclei of the medulla and cord, and exceed some of them, and that the same statement applies to the cells as a whole. I make this statement in view of the recent communication of Dr. Mason before the American Neurological Association, though I do not claim to make it on the same basis of careful and extensive micrometric observations that his communication was based on, but on a general impression derived from repeated examinations which I think are sufficient to determine palpable differences.

I desire the present lines to be looked on as a preliminary communication, and trust to be able before long to submit a more exhaustive and illustrated record of this interesting and suggestive piece of cerebral anatomy to the reader.